arXiv:1203.3278v2 [stat.ME] 21 Mar 2012

On Identity Tests for High Dimensional Data Using RMT

Cheng Wang[a,b], Jing Yang[a], Baiqi Miao[a]

[a] *Department of Statistics and Finance, University of Science and Technology of China, Hefei, Anhui 230026, China*
[b] *wwcc@mail.ustc.edu.cn*

Abstract

In this work, we redefined two important statistics CLRT (Bai et.al., Ann. Stat. **37** (2009) 3822-3840) and LW test(Ledoit and Wolf, Ann. Stat. **30** (2002) 1081-1102) on identity tests for high dimensional data using classic sample covariances. Compared with existing CLRT, the proposed CLRT can accommodate data which have non-zero means and non-Gaussian distributions and the new LW also can be applied to non-Gaussian data set. Simulations demonstrated that the tests had good properties in sizes and powers. What's more, we found CLRT is more sensitive to eigenvalues less than 1 while LW test has more advantages on detecting eigenvalues larger than 1.

Keywords: High dimensional data, Identity test, Random matrix theory
2000 MSC: 62H15, 62H10

---

1. Introduction

Suppose $X_1, \cdots, X_n$ are independent and identically distributed (i.i.d.) p-dimensional random vectors with population covariance matrix Σ_p and our interest is to test

$$H_0: \ \Sigma_p = I_p \ vs. \ H_1: \ \Sigma_p \neq I_p \tag{1.1}$$

where I_p denotes the $p-$dimensional identity matrix. Note that the identity matrix in (1.1) can be replaced by any other positive definite matrix Σ_0 through multiplying the data by $\Sigma_0^{-1/2}$.

In this work, we assume $y_n = p/n \to y \in (0, \infty)$. For canonical statistical analysis, that is the sample size n tends to infinity while the dimension p

remains fixed, one can refer to [1]. When $y < 1$, [3] proposed a correction to the classic likelihood ration test (CLRT) and derived the central limit theorem(CLT) using random matrix theory(RMT). When $y \geq 1$, CLRT is degenerate since the sample covariance is no longer invertible with probability one and [10] gave a new statistics (LW test) which could accommodate situations for any $y > 0$. We note that LW test has drawn much attention in literature which include [16] which considered the test for part of the eigenvalues of Σ_p are equal and [7] which extended LW test to cases $y = 0$ and ∞. There have been also a substantial set of research works motivated by LW test such as [9, 18] and [11].

However, most of these results included CLRT and LW test were derived under Gaussian assumptions or equivalent conditions such as fourth moment equals three. The difficulty to relax Gaussian assumptions is due to the central limit theorems for the linear spectral statistics defined by eigenvalues. More details can be found in [5] which built the CLT for linear spectral statistics of large-dimensional sample covariance matrices under the assumptions on fourth moments and [15] improved the results for general finite fourth moments. The sample covariance matrix defined in [5, 15] is a simplified version of classic sample covariance matrices and the data must had zero mean. Recently, [14] derived the CLT for linear spectral statistics of classic large-dimensional sample covariance matrices. Some other important results include [6, 12] and so on.

Besides Gaussian assumptions, CLRT in [3] actually is only applicable for data with zero-means which is too strict to applications. In this work, we redefined CLRT and LW test using classic sample covariance matrices and derived the CLTs of the two tests in general conditions which can accommodate data which have non-zero means and non-Gaussian distributions. Simulations demonstrate that the proposed tests have good properties in sizes and powers. What's more, compared with LW test, CLRT has its own advantages on detecting the eigenvalues of Σ_p near zero. That is CLRT is more sensitive to eigenvalues less than 1 while LW test has more advantages on detecting eigenvalues larger than 1. In literature, there is also some work not based on sample covariance matrices such as [8] which proposed a test by constructing estimators from the data directly. We also conduct simulations to compare our proposed tests with the one in [8].

The paper is structured as follows. Section 2 introduces the basic data structure and establishes the asymptotic normality of the CLRT and LW test. Section 3 reports simulation studies. All the technical details include proofs and preliminary results in RMT are deferred to the Appendix.

2. Main Results

We assume the observations $X_1, \cdots, X_n$ satisfies a multivariate model ([4])

$$X_i = \Sigma_p^{1/2} Y_i + \mu, \ \ for \ i = 1, \cdots, n \tag{2.2}$$

where μ is a p-dimensional constant vector and the entries of $\mathcal{Y}_n = (Y_{ij})_{p\times n} = (Y_1, \cdots, Y_n)$ are i.i.d. with $EY_{ij} = 0$, $EY_{ij}^2 = 1$ and $EY_{ij}^4 = 3 + \Delta$. Here we introduces two versions of the sample covariance matrices. The classic one is defined as

$$S_n = \frac{1}{n-1} \sum_{k=1}^{n} (X_k - \bar{X})(X_k - \bar{X})',$$

where $\bar{X} = \frac{1}{n} \sum_{k=1}^{n} X_k$ and a simplified version has the form

$$B_n = \frac{1}{n} \sum_{k=1}^{n} (X_k - \mu)(X_k - \mu)'.$$

We refer to [2] and [14] for the differences between S_n and B_n in RMT. Then we can introduce CLRT in [3]

$$\hat{L}_n = \frac{1}{p} tr(B_n) - \frac{1}{p} \log |B_n| - 1 \tag{2.3}$$

where tr denotes the trace.

When $X_i \sim N_p(0, I_p)$ (or $X_i \sim N_p(\mu, I_p)$ where μ is known), [3] derived the CLT of CLRT

$$p(\hat{L}_n - (1 + (1/y_n - 1)\log(1 - y_n))) \xrightarrow{D} \mathbb{N}(-\log(1-y)/2, -2y - 2\log(1-y)).$$

where $\xrightarrow{D}$ denotes convergence in distribution and $\mathbb{N}$ the normal distribution.

When p is larger than sample size n, since the sample covariance matrix S_n is singular, [10] proposed LW test which is defined as

$$\hat{W}_n = \frac{1}{p} tr(S_n - I_p)^2 - \frac{p}{n-1} (\frac{1}{p} tr(S_n))^2 \tag{2.4}$$

If $X_i \sim N_p(\mu, I_p)$ and under some other assumptions, [10] has proven

$$n\hat{W}_n \xrightarrow{D} \mathbb{N}(1, 4). \tag{2.5}$$

In our work, we will redefine the CLRT as

$$L_n = \frac{1}{p} tr(S_n) - \frac{1}{p} \log |S_n| - 1 \tag{2.6}$$

and LW test as

$$W_n = \frac{1}{p} tr(S_n - I_p)^2 - \frac{p}{n-1} (\frac{1}{p} tr(S_n))^2 - \frac{(1+\Delta)(n-2)(n-1)-2}{n(n-1)^2}. \tag{2.7}$$

In (2.6) and (2.7), noting that the statistics are defined by S_n which is invariant under the shift transformation $X_i = X_i + c$, we can assume $\mu = 0$ in (2.2) without loss of generality. By [5], we know almost surely

$$\frac{1}{p} \log |S_n| - \frac{1}{p} \log |\Sigma_p| \xrightarrow{a.s.} -1 - (1/y - 1) \log (1-y)) \equiv d(y).$$

which means $L_n + d(y)$ is a estimator of $(tr(\Sigma_p) - \log |\Sigma_p| - p)/p$. Similarly, W_n is a estimator of $tr(\Sigma_p - I_p)^2/p$. Noting that

$$\Sigma_p = I_p \Leftrightarrow (tr(\Sigma_p) - \log |\Sigma_p| - p)/p = 0 \Leftrightarrow tr(\Sigma_p - I_p)^2/p = 0,$$

therefore, L_n and W_n can act as the statistics to test (1.1) theoretically. Next, we will establish the asymptotic normalities of L_n and W_n.

Theorem 2.1. *When $\Sigma_p = I_p$ and $p/y \to y \in (0, 1)$,*

$$p(L_n - (1 + (1/y_n - 1) \log (1 - y_n))) \xrightarrow{D} \mathbb{N}(m, v)$$

where $m = y(\Delta/2 - 1) - 3 \log(1-y)/2$ and $v = -2y - 2 \log (1-y)$.

Compared with [3] which is only applicable to Gaussian data with known means, Theorem 2.1 was built under general conditions and can be applied to Gaussian data directly. If the population mean is unknown, the CLRT in [3] will behaves poorly and the new one are still applicable. Simulations will be conducted in subsection 3.1.

Theorem 2.2. *Under H_0 and $p/n \to y \in (0, \infty)$,*

$$pW_n \xrightarrow{D} \mathbb{N}(0, 4y^2).$$

Due to the test in [8] and LW test are both based on the estimators of $tr(\Sigma_p - I_p)^2/p$, following are remarks about these two tests and simulations will be shown in next section.

Remark **2.1.** *Direct calculations can show that when $\Sigma = I_p$, W_n is a unbiased estimator of $tr(\Sigma_p - I_p)^2$. In [8], authors also proposed a unbiased estimator V_n of $tr(\Sigma_p - I_p)^2$. What's more, under the conditions of Theorem 2.2,*

$$Var(W_n) \to 4y^2 \quad and \quad Var(V_n) \to 4y^2.$$

However, for Gaussian data, due to W_n is based on S_n which is a complete sufficient statistic for Σ_p, we can claim W_n is the unique best unbiased estimator of $tr(\Sigma_p - I_p)^2$ and $Var(W_n) \le Var(V_n)$ by Lehmann-Scheffé theorem. From simulations, we know $4y^2 \le Var(W_n) \le Var(V_n) \to 4y^2$ and therefore the sizes of our new LW test will behaves better than the test in [8]. For non-Gaussian data such as Gamma and Uniform variables, simulations in subsection 3.2 can show that LW test performs favorably in comparison to existing methods.

Remark **2.2.** *In [8], for comparison purposes, authors assumed $\Delta = 0$ even for gamma random vectors and here from Theorem 2.2, we know it is not reasonable. From Theorem 2.2, when $\Sigma_p = I_p$ and $Y_{i,j} \sim Gamma[4, 0.5]$, by Slutsky's theorem we know*

$$(n\hat{W}_n - 1)/2 \xrightarrow{D} \mathbb{N}(0.75, 1).$$

However, in [8], authors still thought $(n\hat{W}_n - 1)/2 \xrightarrow{D} \mathbb{N}(0, 1)$. Moreover, since $P(Z > \Phi^{-1}(0.95)) = 0.185$ where $Z \sim \mathbb{N}(0.75, 1)$ and Φ is the distribution function of standard norm variables, this explain why the sizes of LW test in [8] are near 0.185 not 5%.

In Theorem 2.2, when $X_i \sim N(\mu, I_p)$ that is $\Delta = 0$, we can get

$$p(\hat{W}_n - \frac{(n-2)(n-1)-2}{n(n-1)^2}) \xrightarrow{D} \mathbb{N}(0, 4y^2)$$

which is in accordance with (2.5) by Slutsky's theorem. By Theorem 2.1 and 2.2, the results depend on Δ. When $EY_{ij} \neq 0$, Theorem 2.1 and 2.2 are still applicable under new assumptions $E(Y_{ij} - EY_{ij})^2 = 1$ and $E(Y_{ij} - EY_{ij})^4 = 3 + \Delta$ by [14].

3. Simulations

We report results from simulation studies which were designed to evaluate the performance of the proposed identity tests for the covariance matrix. For comparison purposes, We also conducted the test proposed by [8](CZZ Test). Here, the ratio y could be estimated by $y_n = p/n$ and three scenarios were considered with respect to the innovation random matrix $(Y_{i,j})_{p\times n}$
(I) $Y_{i,j}$ *i.i.d.* $\sim N(1,1)$ where $\Delta = 0$;
(II) $Y_{i,j}$ *i.i.d.* $\sim Gamma[4, 0.5]$ where $\Delta = 1.5$;
(III) $Y_{i,j}$ *i.i.d.* $\sim Uniform[0, 2\sqrt{3}]$ where $\Delta = -1.2$.

To evaluate the power of the tests, two different population covariance matrices were considered in the simulations. We set $\Sigma_p^{(1)} = diag(1.5\, I_{[0.2p]}, I_{p-[0.2p]})$ or $\Sigma_p^{(2)} = diag(0.5\, I_{[0.2p]}, I_{p-[0.2p]})$, where $[x]$ denoted the integer truncation of x. It is that the diagonal covariance Σ_p had respectively 20% of its diagonal elements being 1.5 or 0.5 whereas the rest were 1.

3.1. CLRT Tests

Firstly, we will study our redefined CLRT and existing CLRT. Since the existing CLRT in [3] only can deal with Gaussian variables with known mean, in our simulations, for each (p, n), we run 1,000 independent replications with real Gaussian variables with means zero. Table 1 reported empirical size and power of our redefined CLRT and existing CLRT for Gaussian variables $Y_{ij} \sim \mathbb{N}(\mu, 1)$ (μ is known and without loss of generality, we assume $\mu = 0$ here). The nominal test level is set to be 5%.

From Table 1, we know even for Gaussian variables with known mean, our CLRT is comparable to one in [3]. As p and n both are increased, the size or power of the two tests are quite close to 5% or 1 and make no much difference. What's more, for the same sample size n, when y gets smaller, the sizes are more close to 5% and the powers are more close to 1. A possible explanation is that if we only have n samples, when p gets smaller (that is y gets smaller), our redefined CLRT or existing CLRT, as the estimator of $(tr(\Sigma_p) - \log|\Sigma_p| - p)/p$, will become more accurate.

Table 1: Performances of redefined CLRT and existing CLRT

	Redefined CLRT			Existing CLRT		
n	$y = 0.25$	$y = 0.5$	$y = 0.75$	$y = 0.25$	$y = 0.5$	$y = 0.75$
$\Sigma_p = I_p,\ Y_{ij} \sim \mathbb{N}(0,1)$						
40	0.077	0.072	0.076	0.071	0.061	0.062
80	0.062	0.061	0.062	0.060	0.056	0.055
160	0.054	0.053	0.054	0.053	0.052	0.053
$\Sigma_p = \Sigma_p^{(1)},\ \ Y_{ij} \sim \mathbb{N}(0,1)$						
40	0.220	0.182	0.177	0.214	0.168	0.162
80	0.397	0.342	0.281	0.397	0.337	0.275
160	0.819	0.769	0.632	0.816	0.762	0.625
200	0.926	0.889	0.815	0.925	0.895	0.816
$\Sigma_p = \Sigma_p^{(2)},\ \ Y_{ij} \sim \mathbb{N}(0,1)$						
40	0.371	0.369	0.272	0.370	0.367	0.278
80	0.841	0.749	0.618	0.840	0.762	0.641
160	1	1	0.986	1	0.999	0.990
$\Sigma_p = I_p,\ \ Y_{ij} \sim \mathbb{N}(1/4,1)$						
100	0.066	0.051	0.059	0.689	0.837	0.834

If the true mean μ is not zero and we still use the CLRT in [3], the results of the last part in Table 1 is the experiment for $Y_{ij} \sim \mathbb{N}(1/4, 1)$. It can be found that the existing CLRT behaves very poorly and our CLRT is still applicable.

3.2. LW test VS CZZ test

From Theorem 2.2, we know the CLT of LW test depends on Δ and it is not reasonable that in [8] authors assumed $\Delta = 0$ even for gamma random vectors. Here we will repeat part of the simulations in [8] and the nominal significant level of the tests was 5%. Simulation results are reported in Table 2 where the performances for Gaussian data are based on 10^4 simulations and for non-Gaussian data results are based on 10^3 simulations.

Obviously, the proposed LW test is not so bad as shown in Table 1 of [8]. We observe from Table 2 that compared with CZZ test, LW test has a better empirical size for Gaussian data which were assuring in Remark 2.1. For Gamma or Uniform variables, the sizes of LW test and CZZ test are comparable and when p or n increases, the sizes both tend to the nominal 5% level. It is noted that Table 1 in [8] was the results for sphericity test, not

Table 2: Performances of LW test and CZZ test

	LW test				CZZ test			
p	$n = 20$	$n = 40$	$y = 60$	$n = 80$	$n = 20$	$n = 40$	$n = 60$	$n = 80$
Gaussian random vectors								
38	0.0659	0.0569	0.0601	0.0551	0.0730	0.0573	0.0637	0.0576
55	0.0693	0.0611	0.0593	0.0537	0.0732	0.0616	0.0600	0.0532
89	0.0650	0.0558	0.0603	0.0531	0.0692	0.0585	0.0620	0.0550
159	0.0635	0.0600	0.0608	0.0564	0.0673	0.0609	0.0598	0.0569
Gamma random vectors								
38	0.096	0.079	0.069	0.065	0.078	0.066	0.064	0.058
55	0.092	0.089	0.085	0.087	0.069	0.076	0.070	0.089
89	0.101	0.089	0.070	0.068	0.086	0.075	0.053	0.067
159	0.086	0.065	0.061	0.062	0.075	0.051	0.055	0.054
Uniform random vectors								
38	0.051	0.044	0.046	0.051	0.059	0.050	0.049	0.056
55	0.064	0.058	0.056	0.043	0.073	0.064	0.059	0.044
89	0.057	0.049	0.056	0.043	0.069	0.056	0.059	0.047
159	0.050	0.051	0.056	0.056	0.061	0.056	0.059	0.062

identity test. Since authors claimed the simulation results for the identity test followed very much similar patterns to those of the sphericity test, here for comparison purposes, we still use Table 1 in [8] for identity test.

Above all, the performances of the proposed LW test are comparable to CZZ test and for Gaussian data, the new LW test even has a better size. Compared with CZZ test which does not depend on Δ, LW test has its own advantages on computational cost. Actually, in order to compute LW test, we only need the sample covariance matrix S_n while CZZ test is consist of five parts.

3.3. Powers of LW and CLRT tests

Results were reported in Figure 1, corresponding to innovation random matrices (I), (II) and (III) and $\Sigma_p = \Sigma_p^{(1)}$ or $\Sigma_p = \Sigma_p^{(2)}$. The power results in Figure 1 showed CLRT and LW tests both approached to 1 when n was increased. For $\Sigma_p^{(1)}$, that is part of the eigenvalues of Σ_p were larger than 1, the power of CLRT was a little worse than the one of LW test and for $\Sigma_p^{(2)}$, when part of eigenvalues were less than 1, CLRT behaved better than LW test. The reason is due to the differences between $tr(\Sigma_p) - \log|\Sigma_p| - p$ and



Figure 1: Realized powers of LW test and CLRT for different sample size n and $p/n = 0.5$. The top three are for $\Sigma_p = \Sigma_p^{(1)}$ and the bottom three are for $\Sigma_p = \Sigma_p^{(2)}$.

$tr(\Sigma_p - I_p)^2$ which offer avenues for further research and we would not pursue it here due to a limited space.

4. Conclusions

In this work, we redefined two identity tests CLRT and LW test for high dimensional data. Compared with existing CLRT, the new CLRT and LW test can accommodate data which have non-zero means and non-Gaussian distributions. Simulations demonstrated that the tests had good properties in sizes and powers. What's more, from the simulations, we found CLRT is not the best one even for Gaussian variables although CLRT came from the likelihood functions of normal distributions. Also, the powers of the tests (included CZZ test) depend on the population covariance matrix and our next work will focus on performances of LW test (or CZZ test) and CLRT theoretically.

Acknowledgements

The authors would like to thank the associate editor and an anonymous referee for their helpful comments. We would especially appreciate the anonymous referee whose insightful suggestions have improved the paper presentation significantly.

5. Appendix

5.1. Preliminary results in RMT

Suppose A_n is an $n \times n$ Hermitian matrix with eigenvalues $\lambda_1, \cdots, \lambda_n$. Define the empirical spectral distribution (ESD) of A_n as

$$F^{A_n}(x) = \frac{1}{n} \sum_{i=1}^{n} I(\lambda_i \leq x).$$

The limit distribution of F^{A_n} is called the limiting spectral distribution (LSD) of the sequence $\{A_n\}$.

And the Stieltjes transform of F^{A_n} is given by

$$m^{F^{A_n}}(z) = \int \frac{1}{x - z} dF^{A_n}(x) = \frac{1}{n} tr(A_n - zI_n)^{-1},$$

where $z = \mu + i\nu \in \mathcal{C}^+$. By the inverse formula,

$$F^{A_n}\{[a, b]\} = \lim_{v \to 0^+} \frac{1}{\pi} \int_a^b Im(m^{F^{A_n}}(x + iv))dx. \tag{5.8}$$

Here we need another sample covariance matrix which is defined as

$$\mathfrak{S}_n = \frac{1}{n} \sum_{k=1}^{n} (X_k - \bar{X})(X_k - \bar{X})' = \frac{n-1}{n} S_n. \tag{5.9}$$

If the spectral norm of Σ_p is bounded by a positive constant and F^{Σ_p} converges weakly to a non-random distribution H as $p \to \infty$, by [17] or [13], with probability 1, $F^{\mathfrak{S}_n}$ and F^{B_n} tend to the same probability distribution $F_{y,H}$, whose Stieltjes transform $m = m(z)$ ($z \in \mathcal{C}^+$) satisfies

$$m = \int \frac{1}{t(1 - y - yzm) - z} dH(t). \tag{5.10}$$

Denoting $G_n(x) = p(F^{\mathfrak{S}_n}(x) - F_{y_n, H_n}(x))$, for any analytic function f, $\int f(x)dG_n(x)$ converges weakly to a Gaussian variable X_f under some assumptions on Σ_p by [14].

When $\Sigma_p = I_p$, $F_{y,H}$ is standard MP law F_y whose density function is

$$g_y(x) = \frac{1}{2\pi yx} \sqrt{((1 + \sqrt{y})^2 - x)(x - (1 - \sqrt{y})^2))},\ (1 - \sqrt{y})^2 \leq x \leq (1 + \sqrt{y})^2$$

and from (5.10), we know

$$m = \frac{1}{1 - y - yzm - z}.$$

Writing $\underline{m} = ym - \frac{1-y}{z}$, by [14], we have

$$\begin{aligned} EX_f &= -\frac{1}{2\pi i}\int \frac{y\underline{m} f(-\frac{1}{\underline{m}} + \frac{y}{1+\underline{m}})}{(1+\underline{m})((1+\underline{m})^2 - c\underline{m}^2)} d\underline{m} - \frac{\Delta}{2\pi i}\int \frac{y\underline{m} f(-\frac{1}{\underline{m}} + \frac{y}{1+\underline{m}})}{(1+\underline{m})^3} d\underline{m} \\ &\quad + \frac{y}{2\pi i}\int \frac{f(-\frac{1}{\underline{m}} + \frac{y}{1+\underline{m}})}{(1+\underline{m})(y\underline{m} - 1 - \underline{m})} d\underline{m}. \end{aligned} \tag{5.11}$$

and

$$\begin{aligned} Var(X_f) &= -\frac{1}{2\pi^2}\int\int f(z_1)f(z_2)\frac{1}{(\underline{m}(z_1) - \underline{m}(z_2))^2} d\underline{m}(z_1) d\underline{m}(z_2) \\ &\quad - \frac{y\Delta}{4\pi^2}\left(\int \frac{f(z)}{(1+\underline{m}(z))^2} d\underline{m}(z)\right)^2. \end{aligned} \tag{5.12}$$

The contours in (5.11) and (5.12) are both contained in the analytic region for the function f and both enclose the support of $F_{y_n,H_n}(x)$ for large n. Moreover, the contours in (5.12) are disjoint.

5.2. Proofs of Theorem 2.1

Writing $f(x) = x - \log(x) - 1$, we have

$$L_n = \int f(x) dF^{S_n}(x)$$

and

$$\begin{aligned} & p(L_n - \int f(x) dF_{y_n}(x)) \\ &= p(\int f(x) dF^{\mathfrak{S}_n}(x) - \int f(x) dF_{y_n}(x)) + p(\int f(x) dF^{S_n}(x) - \int f(x) dF^{\mathfrak{S}_n}(x)) \\ &= p(\int f(x) dF^{\mathfrak{S}_n}(x) - \int f(x) dF_{y_n}(x)) + (tr(S_n) - log|S_n| - tr(\mathfrak{S}_n) + \log|\mathfrak{S}_n|) \\ &= p(\int f(x) dF^{\mathfrak{S}_n}(x) - \int f(x) dF_{y_n}(x)) + \frac{1}{n-1}tr(\mathfrak{S}_n) - p\log(\frac{n}{n-1}) \\ &= p(\int f(x) dF^{\mathfrak{S}_n}(x) - \int f(x) dF_{y_n}(x)) + o(1). \end{aligned}$$

From [14], $p(\int f(x)dF^{\mathfrak{S}_n}(x)-\int f(x)dF_{y_n}(x))$ converges weakly to the Gaussian variable X_f. Next we calculate the mean and variance of X_f. The following results have been given in Bai et.al.(2008)

$$\int f(x)dF_{y_n}(x)=1-\frac{y_n-1}{y_n}\log(1-y_n),$$

$$-\frac{1}{2\pi i}\int\frac{y\underline{m}f(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})}{(1+\underline{m})((1+\underline{m})^2-c\underline{m}^2)}d\underline{m}=-\frac{1}{2}\log(1-y), \tag{5.13}$$

and

$$-\frac{1}{2\pi^2}\int\int\frac{f(z_1)f(z_2)}{(\underline{m}(z_1)-\underline{m}(z_2))^2}d\underline{m}(z_1)d\underline{m}(z_2)=-2y-2\log{(1-y)}. \tag{5.14}$$

Also, from [15], we know

$$\frac{1}{2\pi i}\oint(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}}-1)\frac{y\underline{m}}{(1+\underline{m})^3}d\underline{m}=0.$$

Therefore, to get EX_g, we still need to calculate

$$\begin{aligned}
& \frac{y}{2\pi i}\oint\log(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})\frac{\underline{m}}{(1+\underline{m})^3}d\underline{m} \\
= & \frac{y}{2\pi i}\oint\log(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})d(-\frac{1}{1+\underline{m}}+\frac{1}{2(1+\underline{m})^2}) \\
= & \frac{y}{2\pi i}\oint\frac{\frac{1}{\underline{m}^2}-\frac{y}{(1+\underline{m})^2}}{-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}}}(\frac{1}{1+\underline{m}}-\frac{1}{2(1+\underline{m})^2})d\underline{m} \\
= & \frac{y}{2\pi i}\oint(\frac{1+\underline{m}}{\underline{m}(y\underline{m}-\underline{m}-1)}-\frac{y\underline{m}}{(\underline{m}+1)(y\underline{m}-\underline{m}-1)})(\frac{1}{1+\underline{m}}-\frac{1}{2(1+\underline{m})^2})d\underline{m} \\
= & \frac{y}{2\pi i}\oint(\frac{1}{\underline{m}(y\underline{m}-\underline{m}-1)}-\frac{1}{2\underline{m}(\underline{m}+1)(y\underline{m}-\underline{m}-1)})d\underline{m} \\
= & \frac{y}{2},
\end{aligned} \tag{5.15}$$

and

$$
\begin{aligned}
& \frac{y}{2\pi i}\int \frac{f(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})}{(1+\underline{m})(y\underline{m}-1-\underline{m})}d\underline{m} \\
= \ & \frac{y}{2\pi i}\int \frac{-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}}-1-\log(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})}{(1+\underline{m})(y\underline{m}-1-\underline{m})}d\underline{m} \\
= \ & -y+\frac{1}{2\pi i}\int \log(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})(\frac{1}{1+\underline{m}}-\frac{1}{\underline{m}-\frac{1}{y-1}})d\underline{m} \\
= \ & -y-\log(1-y) \qquad (5.16)
\end{aligned}
$$

where we used the following results

$$\oint \frac{1}{(\underline{m}+1)^k(y\underline{m}-\underline{m}-1)}d\underline{m}=0,\ \ k=1,2,3,$$

and one equality in [5]

$$\frac{1}{\pi i}\int \log(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})(\frac{1}{1+\underline{m}}-\frac{1}{\underline{m}-\frac{1}{y-1}})d\underline{m}=-2\log(1-y).$$

By (5.13), (5.15) and (5.16), $EX_f = y(\Delta/2-1)-3\log(1-y)/2$.

Through a routine calculation, we have

$$\frac{1}{2\pi i}\oint f(-\frac{1}{\underline{m}}+\frac{y}{1+\underline{m}})\frac{1}{(1+\underline{m})^2}d\underline{m}=0 \qquad (5.17)$$

By (5.14) and (5.17), we have $Var(X_f)=-2y-2\log(1-y)$.

The proof of Theorem 2.1 is complete.

5.3. Proofs of Theorem 2.2

Noticing that $p(\frac{1}{p}tr(\mathfrak{S}_n)-1)$ satisfies CLT and $\frac{1}{p}tr(\mathfrak{S}_n)\to 1, a.s.$, we have

$$p((\frac{1}{p}tr(\mathfrak{S}_n))^2-2\frac{1}{p}tr(\mathfrak{S}_n)+1)=p([\frac{1}{p}tr(\mathfrak{S}_n)]-1)^2=o(1).$$

By (5.9), we have

$$
\begin{aligned}
& p[\hat{W}_n - (\frac{1}{p}tr(\mathfrak{S}_n - I_p)^2 - \frac{2}{n}tr(\mathfrak{S}_n) + \frac{p}{n})) \\
& = tr(\frac{n}{n-1}\mathfrak{S}_n - I_p)^2 - tr(\mathfrak{S}_n - I_p)^2 + (\frac{1}{n} - \frac{n^2}{(n-1)^3})(tr(\mathfrak{S}_n))^2 + o(1) \\
& = (\frac{n^2}{(n-1)^2} - 1)tr(\mathfrak{S}_n^2) - 2(\frac{n}{n-1} - 1)tr(\mathfrak{S}_n) + (\frac{1}{n} - \frac{n^2}{(n-1)^3})(tr(\mathfrak{S}_n))^2 \\
& = -y^2 + o(1)
\end{aligned}
$$

and

$$
\begin{aligned}
& p(\frac{1}{p}tr(\mathfrak{S}_n - I_p)^2 - \frac{2}{n}tr(\mathfrak{S}_n) + \frac{p}{n}) - \int((x-1)^2 - 2yx + y)dp(F^{\mathfrak{S}_n}(x) - F_{y_n}(x)) \\
& = (y - y_n)(2tr(\mathfrak{S}_n) - p) + p\int((x-1)^2 - 2yx + y)dF_{y_n}(x) \\
& = 2(y - y_n)(tr(\mathfrak{S}_n) - p) \\
& = o(1)
\end{aligned}
$$

Writing $g(x) = (x-1)^2 - 2yx + y$, by [14], $\int g(x)dp(F^{\mathfrak{S}_n}(x) - F_{y_n}(x))$ converges weakly to gaussian variable X_g and through a routine calculation

$$EX_g = y + y\Delta + y^2 = y(1 + \Delta + y).$$

To get the variance $Var(X_g)$, we need

$$
\begin{aligned}
& \oint \frac{g(z(m_1))}{(m_1 - m_2)^2}dm_1 \\
= & \oint \frac{1}{(m_1 - m_2)^2}[(-\frac{1}{m_1} + \frac{y}{1+m_1} - 1)^2 + 2y(-\frac{1}{m_1} + \frac{y}{1+m_1} + 1)]dm_1 \\
= & \frac{4\pi i y^2}{(m_2+1)^3} - \frac{4\pi i y^2}{(m_2+1)^2} = -\frac{4\pi i y^2 m_2}{(m_2+1)^3},
\end{aligned}
$$

$$-\frac{2y^2}{\pi i}\oint g(z(m))\frac{m}{(m+1)^3}dm = 4y^2,$$

and

$$\frac{1}{2\pi i}\int g(-\frac{1}{m} + \frac{y}{1+m})\frac{1}{(1+m)^2}dm = 0$$

Then

$$Var(X_g) = 4y^2.$$

Above all, we can get

$$p\hat{W}_n \xrightarrow{D} \mathbb{N}(y, 4y^2).$$

By Slutsky's theorem, the proof of Theorem 2.2 is complete.

References

[1] T.W. Anderson, An Introduction to Multivariate Statistical Analysis. Hoboken, NJ: Wiley (2003).

[2] Z.D. Bai, and J.W. Silverstein, Spectral Analysis of Large Dimensional Random Matrices, Beijing: Scientific Press (2006).

[3] Z.D. Bai, D.D. Jiang, J.F. Yao, and S.R. Zheng, Corrections to Lrt on Large-Dimensional Covariance Matrix by Rmt. Annals of Statistics 37 (2009) 3822-3840.

[4] Z.D. Bai, and H. Saranadasa, Effect of high dimension: By an example of a two sample problem. Statistica Sinica 6 (1996) 311-329.

[5] Z.D. Bai, and J.W. Silverstein, CLT for linear spectral statistics of large-dimensional sample covariance matrices. Annals of Probability 32 (2004) 553-605.

[6] Z.D. Bai, X.Y. Wang, and W. Zhou, Functional CLT for sample covariance matrices. Bernoulli 16 (2010) 1086-1113.

[7] M. Birke, and H. Dette, A note on testing the covariance matrix for large dimension. Statistics and Probability Letters 74 (2005) 281-289.

[8] S.X. Chen, L.X. Zhang, and P.S. Zhong, Tests for High-Dimensional Covariance Matrices. Journal of the American Statistical Association 105 (2010) 810-819.

[9] T.J. Fisher, X.G. Sun, and C.M. Gallagher, A new test for sphericity of the covariance matrix for high dimensional data. Journal of Multivariate Analysis 101 (2010) 2554-2570.

[10] O. Ledoit, and M. Wolf, Some hypothesis tests for the covariance matrix when the dimension is large compared to the sample size. Annals of Statistics 30 (2002) 1081-1102.

[11] Z.Y. Lin, and Y.B. Xiang, A hypothesis test for independence of sets of variates in high dimensions. Statistics and Probability Letters 78 (2008) 2939-2946.

[12] A. Lytova, and L. Pastur, Central Limit Theorem for Linear Eigenvalue Statistics of Random Matrices with Independent Entries. Annals of Probability 37 (2009) 1778-1840.

[13] G.M. Pan, Strong convergence of the empirical distribution of eigenvalues of sample covariance matrices with a perturbation matrix. Journal of Multivariate Analysis 101 (2010) 1330-1338.

[14] G.M. Pan, Comparison between two types of sample covariance matrices, manuscript (2011).

[15] G.M. Pan, and W. Zhou, Central limit theorem for signal-to-interference ratio of reduced rank linear receiver. Annals of Applied Probability 18 (2008) 1232-1270.

[16] J.R. Schott, A high-dimensional test for the equality of the smallest eigenvalues of a covariance matrix. Journal of Multivariate Analysis 97 (2006) 827-843.

[17] J.W. Silverstein, and Z.D. Bai, On the Empirical Distribution of Eigenvalues of a Class of Large Dimensional Random Matrices. Journal of Multivariate Analysis 54 (1995) 175-192.

[18] M.S. Srivastava, Some tests concerning the covariance matrix in high dimensional data. J. Japan Statist. Soc. 35 (2005).